


BONE AND ARTICULAR CARTILAGE POLISH NATIONAL OF LIMBS CANCER (C40-C41) - 2019


Factsheet


	Morbidity		
	Males	Females	Both
Number of cases in 2019	158	157	315
Crude rate	0.9	0.8	0.8
Standardized rate (ASE)	0.9	0.8	0.8
Cumulative risk (to age 74)	1 of 153447	1 of 176569	-
Percentage of all cancers	0.2%	0.2%	0.2%
Predicted number of cases in 2024	158	135	293
Predicted standardized rate (ASE) in 2024	0.9	0.7	0.8













Males Females Both Number of cases in 2019 145 101 246 Crude rate 0.8 0.5 0.6 Standardized rate (ASE) 0.9 0.5 0.7 Cumulative risk (to age 74) 1 of 203021 1 of 334604 Percentage of all cancers 0.3% 0.2% 0.2% Predicted number of cases in 2024 126 226 100 Predicted standardized rate (ASE) in 2024 0.8 0.5 0.6







20-34 15.2%


0-19 18.4%







Deaths, females





BONE AND ARTICULAR CARTILAGE OF LIMBS **CANCER - 2019**



Factsheet

Incidence by sex









Standardized rate (ASE)

Standardized rate (ASE)



09 2011 2013 2015 2017 2019 2010 2012 2014 2016 2018 2002 2004 2006 2008

Incidence by sex (standardized rate ASE)



Male Female

Bone and articular cartilage of limbs cancer risk factors:

- age the risk of developing the disease is greater in adolescence, •
- gender osteosarcoma is more common in men, .
- height children with osteosarcoma tend to be taller than their peers.



1.00



©2022 Polish National Cancer Registry

Standardized rate (ASE)

Mortality

0.8

0.8

0.7

0.3

0.5

0.6

0.6

0.6

0.7

0.7

0.8

0.8

0.8

0.50

0.6

1.0